PACKAGE LEAFLET

For animal treatment only

TIAMUVET®

medicated premix 2% of tiamulin for pigs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing Authorisation Holder:

PROVET S.A.

120 Eleftherias Avenue, Eleousa, Zitsa, 45500 Ioannina, Greece

Tel.: +30 2105508500, +30 2105575770-3

E-mail: vet@provet.gr

Manufacturer responsible for batch release:

PROVET S.A. (FACTORY)

Nikiforou Foka & Ag.Anargyron, Thesi Vrago, 193 00 Aspropyrgos, Attiki, Greece

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

TIAMUVET

Medicated premix 2%

Tiamulin

for pigs

3. STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS

100 g of product contain:

Active substance: 2.5g Tiamulin Hydrogen Fumarate 80% (equivalent to 2g Tiamulin Hydrogen

Fumarate)

Excipient: Maize starch

4. INDICATIONS

TIAMUVET® premix 2% is indicated for the prevention and treatment of mycoplasma infections of pigs (enzootic pneumonia, pleuropneumonia, dysentery and leptospirosis).

5. CONTRAINDICATIONS

Do not administer to boars or pregnant sows except for the period 4 weeks prior and 4 weeks after parturition.

6. ADVERSE REACTIONS

The use of antimicrobial therapy could lead to growth of non-sensitive bacteria and fungi. Skin rash may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Pigs

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Administer TIAMUVET® premix 2% per os . It is mixed in the final feed at the following dosage rates:

	Enzootic pneumonia	Dysentery	Leptospirosis
PREVENTION	200 ppm active substance or 10kg of TIAMUVET/ tons for 5 days and then 2kg of product/ tons for 6 weeks	40 ppm active substance or 2kg of product/ tons for 2-3 weeks in periods of increased stress (weaning-grouping)	
TREATMENT	200 ppm active substance or 10kg of product/ tons for 10 days and then 2 kg of product/ tons for 2-3 weeks	100-150 ppm active substance 5-7.5 kg of product/ tons for 7-10 days and then 2kg of product/ tons for 2-3 weeks	200 ppm active substance or 10kg of product/ tons for 10 days

9. ADVICE ON CORRECT ADMINISTRATION

Do not administer concurrently with ionophores (polyether), such as monensin, salinomycin, maduramicin, naracin as severe growth despression or death may result.

10. WITHDRAWAL PERIOD

Meat: 14 days

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store below 25°C,

Protected from light.

Shelf life after incorporation into the feed: 1 month

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

12. SPECIAL WARNINGS

The use of antimicrobial therapy could lead to growth of non-sensitive bacteria and fungi. Should there be no response to treatment within 5 days, re-evaluate the diagnosis. When mixing, avoid direct contact of TIAMUVET® premix 2% with the skin and mucous membranes.

13.SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

FEBRUARY 2008

15. OTHER INFORMATION

PACKAGES

Box of 5 kg

Sack of 20 kg and 25 kg

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorization holder.

under veterinary medical prescription
KEEP OUT OF THE SIGHT AND REACH OF CHILDREN