

LEAFLET

**NEOMYVET<sup>®</sup> w.s.p. 70%**  
**FOR ANIMAL TREATMENT ONLY**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORIZATION HOLDER AND OF THE MANUFACTURING AUTHORIZATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorization holder:

**PROVET S.A.**

77, Posidonos Avenue

174 55 Alimos, Attiki

Tel.: +30 2105575770-3, +30 210 5508500

Fax: +30 2105575830

E-mail: vet@provet.gr

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

**NEOMYVET<sup>®</sup> w.s.p.70%**

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS**

1 g of product contains:

**Active substance:** 700 mg Neomycin sulfate (equivalent to 490 mg Neomycin base)

**Excipient:** Sucrose powder

**4. INDICATION(S)**

NEOMYVET<sup>®</sup> w.s.p. 70% is recommended for the prevention and treatment of pigs and poultry bacterial enteritis, caused by microorganisms susceptible to neomycin.

**5. CONTRAINDICATIONS**

None known

**6. ADVERSE REACTIONS**

None known

If you notice any serious effects or other effects not mentioned on this package, please inform your veterinary surgeon.

**7. TARGET SPECIES**

Pigs, Poultry

**8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

The oral administration of daily dose of neomycin sulfate for pigs and poultry is 11 mg/ kg of b.w. This dose is ensured by diluting the product in the drinking water at the following rates:

Dosage

In order to calculate the administered in the drinking water, dose of NEOMYVET<sup>®</sup> w.s.p. 70% for pigs and poultry, one must multiply the kilos of body weight 16 times and divide the product by 1000. The result expresses the quantity of NEOMYVET<sup>®</sup> w.s.p. 70% in grams that must be dissolved in the water. The correct dose is added in the drinking water and should be consumed by poultry within the next 12 hours. The addition of NEOMYVET<sup>®</sup> w.s.p. 70% in the drinking water must be done daily. During the course of treatment there should not be other water sources available.

Duration of treatment

For the prevention of bacterial enteritis the administration of NEOMYVET<sup>®</sup> w.s.p. 70% is recommended for at least 5 to 10 days, when the danger is more eminent. For the treatment of bacterial enteritis the product is administered for 3-5 days. If there is no improvement within 3 days of treatment, the diagnosis should be re-evaluated.

## **9. ADVICE ON CORRECT ADMINISTRATION**

If there is no improvement within 3 days of treatment, the diagnosis should be re-evaluated and the antibacterial agent altered. Should the diarrhea be complicated by other infectious agents, administer systematic antibacterial treatment.

## **10. WITHDRAWAL PERIOD**

Meat: Pigs:	0 days
Broilers:	1 day
Poultry (turkeys, ducks):	0 days
Eggs:	0 days

## **11. SPECIAL STORAGE PRECAUTIONS**

Store at temperatures below 25 °C.

Shelf life after dilution in the drinking water: 24 hours

## **12. SPECIAL WARNING(S)**

Some feeds contain argillaceous substances, such as bentonite, that can act as a transmitter of ions. This ability of transmitting of ions can have an effect on the bioavailability of aminoglycosides, neomycin included.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

## **14. OTHER INFORMATION**

### **PACKAGE**

Plastic container of 500 g

Marketing Authorization Number: 19947/22-04-2003/K-0147901

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorization holder.

**\*under veterinary medical prescription\***  
**KEEP OUT OF THE REACH AND SIGHT OF CHILDREN**