

LEAFLET

THIABENDAZOLE/ PROVET tablets
FOR ANIMAL TREATMENT ONLY

1. NAME AND ADDRESS OF THE MARKETING AUTHORIZATION HOLDER AND OF THE MANUFACTURING AUTHORIZATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorization holder:

PROVET S.A.

77, Posidonos Avenue

174 55 Alimos, Attiki

Tel.: +30 2105575770-3, +30 210 5508500

Fax: +30 2105575830

E-mail: vet@provet.gr

2. NAME OF THE VETERINARY MEDICINAL PRODUCT
THIABENDAZOLE/ PROVET tablets

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

Each tablet contains:

Active substance: 2 g Thiabendazole (2-(4-Thiazolyl)-benzimidazole)

Excipients: Cellulose microcrystalline, Hydrogenated castor oil, Sodium starch glycollate, Sodium lauryl sulfate, Magnesium stearate, Povidone (Polyvinylpyrrolidone), Lactose, Polysorbate 80, Starch maize, Alcohol

4. INDICATION(S)

It is recommended for the treatment of parasitic infestations of animals caused by mature and immature stages of gastrointestinal and pulmonary roundworms. In severe cases of pulmonary strongylosis, use double the dose indicated for gastrointestinal strongylosis. It is also recommended for the treatment against *hepatic dicrocoeliosis*.

5. CONTRAINDICATIONS

None

6. ADVERSE REACTIONS

Prolonged and improper dosage may lead to the development of resistant parasitic strains especially in *Haemonchus contortus*, *Ostertagia circumcincta* and *Trichostrongylus colubriformis*. This is avoided by using the appropriate therapeutic dose, at regular intervals, and the alteration of the chemical group of anthelmintics.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Cattle, Sheep, Goats

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

It is administered per os by hand or with a balling gun. They can be also crushed and dissolved in water.

ANIMAL SPECIES	BODY WEIGHT	NUMBER OF TABLETS			
		ROUTINE WORMING	SEVERE PARASITISM	LUNGWORMS	DICROCOELIOSIS
Lambs-Kids	under 25 kg	½	1	1	2 ½
Sheep-Goats	over 25 kg	1	1 ½	2	4
Rams-Goat	over 50 kg	1 ½	2 ½	3	5
Cattle	per 30 kg	1	-	2	-

9. ADVICE ON CORRECT ADMINISTRATION

None

10. WITHDRAWAL PERIOD

Meat: 5 days

Milk: 3 days

11. SPECIAL STORAGE PRECAUTIONS

Store in a cool and dry place.

12. SPECIAL WARNING(S)

Thiabendazole is incompatible with products containing copper and cobalt salts. It is synergistic with other anthelmintics and especially with bithionol sulfoxide.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. OTHER INFORMATION

PACKAGE

Box of 50 tablets in blister

Marketing Authorization Number: 8815/28-03-1989/K-0037801

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorization holder.

under veterinary medical prescription
KEEP OUT OF THE REACH AND SIGHT OF CHILDREN